

Dialectics of Knowledge Representation in a Granular Rough Set Theory

A. Mani*

Department of Pure Mathematics
University of Calcutta
9/1B, Jatin Bagchi Road
Kolkata-700029, India
Web: <http://www.logicamani.in>
Email: a.mani.cms@gmail.com

Refereed Paper: ICLA, 9-12th January'2013, Chennai

Abstract. The concepts of rough and definite objects are relatively more determinate than those of granules and granulation in general rough set theory (RST) [1]. Representation of rough objects can however depend on the dialectical relation between granulation and definiteness. In this research, we exactify this aspect in RST over proto-transitive approximation spaces. This approach can be directly extended to many other types of RST. These are used for formulating an extended concept of knowledge interpretation (KI)(relative the situation for classical RST) and the problem of knowledge representation (KR) is solved. These will be of direct interest in granular KR in RST as developed by the present author [2] and of rough objects in general. In [3], these have already been used for five different semantics by the present author.

keywords: Rough Objects, Proto Transitivity, Granulation, RYS, Lattice Theory of Definite Objects, Axiomatic Theory of Granules, Contamination Problem, Granular Knowledge, Knowledge Representation.

1 Introduction

General transitive relations are of much interest in a wide variety of application scenarios including vagueness and preference. But the semantics for the former cases have not been considered in the literature prior to this and [3] by the present author. Here we develop the KI over a granular representation of rough objects. We also show the gaping holes in meaning that would be left by omitting granularity in the considerations.

Rough objects as explained in [4,1] are collections of objects in a classical domain (Meta-C) that appear to be indistinguishable among themselves in another rough semantic domain (Meta-R). But their representation in most RSTs in purely order theoretic terms is not known. In this research paper, we do this for a specific type of RST over proto-transitive approximation spaces (PRAS)

* I would like to thank Prof Mihir Chakraborty for discussions on PRAX.

developed recently by the present author in [3]. The method can be directly extended to many other types of RST. In [3], five different algebraic semantics in slightly different rough semantic domains are also developed by the present author. In general, rough objects correspond to concepts in the Pawlak-sense KI as they correspond to the *objects that can be perceived in the rough semantic domain*. Here we focus on the generalized approach to granular KI in general RST initiated in [2]. Relative the extended KI in a semantic domain placed between the classical (Meta-C) and Meta-R (of rough objects), the problem of KR is solved for reflexive PRAS (PRAX).

From a simplified view, rough objects may be represented by pairs or tuples of definite objects under some conditions. In classical RST, definite objects are precisely those that satisfy $x^l = x^u = x$. Pairs of definite objects of the form (a, b) satisfying $a \subseteq b$ necessarily represent rough objects and every rough object can be so represented. Further intervals of the form $]a, b[= \{c : a \subset c \subset b, \}$ represent rough objects if and only if b covers a . However not all intervals of the form correspond to rough objects without the covering condition. This relation becomes more complicated in more general RSTs as in the case of PRAX. We characterize this in this research.

Classical RST is RST starting from an approximation space of the form $\langle \underline{S}, R \rangle$ with $\underline{S}$ being a set and R being an equivalence relation on $\underline{S}$. *Weak transitivity* of [5] is *proto-transitivity* here and distinct from the usage in [6].

Definition 1. A binary relation R on a set S is said to be weakly-transitive, transitive or proto-transitive respectively on S if and only if S satisfies

- ★ $(\forall x, y, z)(Rxy, Ryz \& x \neq y \neq z \longrightarrow Rxz)$ (i.e. $(R \circ R) \setminus \Delta_S \subseteq R$ (where $\circ$ is relation composition) , or
- ★ $(\forall x, y, z)(Rxy \& Ryz \longrightarrow Rxz)$ (i.e. $(R \circ R) \subseteq R$), or
- ★ $(\forall x, y, z)(Rxy, Ryz, Ryx, Rzy \& x \neq y \neq z \longrightarrow Rxz)$, respectively. Proto-transitivity of R is equivalent to $R \cap R^{-1} = \tau(R)$ being weakly transitive.

$Ref(S)$, $r\tau(S)$, $w\tau(S)$, $p\tau(S)$, $EQ(S)$ will respectively denote the set of reflexive, transitive, weakly transitive, proto transitive, and equivalence relations on the set S respectively.

Proposition 1. In general, $w\tau(S) \subseteq p\tau(S)$. If $R \in Ref(S)$, then $R \in p\tau(S)$ if and only if $\tau(R) \in EQ(S)$.

Different formal definitions of granules have been used in the literature on rough sets and in granular computing. An improved version of the axiomatic theory of granules introduced in [4] can be found in [1] and a formal simplified version in [7]. The axiomatic theory is capable of handling most contexts and is intended to permit relaxation of set-theoretic axioms at a later stage. The axioms are considered in the framework of Rough Y-Systems (RYS) that maybe seen as a generalized form of *abstract approximation spaces* [8] and approximation framework [9]. It includes relation-based RST, cover-based RST and more. These structures are provided with enough structure so that a classical semantic domain (Meta-C) and at least one rough semantic domain (called Meta-R) of

roughly equivalent objects along with admissible operations and predicates are associable. But the exact way of association is not something absolute as there is no real end to recursive approximation processes of objects.

2 Motivation and Examples

Generalized transitive relations occur frequently in general information systems, but are often not recognized as such and there is hope for improved semantics and KI relative the situation for purely reflexive relation based RST. Not all of the definable approximations have been investigated in even closely related structures of general RST. Contamination-free semantics [1] for the contexts are also not known. Finally these relate to RYS and variants. A proper characterization of roughly equal (requal) objects is also motivated by [1].

Example-1:

Let $\S = \{a, b, c, e, f, g, h, l, n\}$ and let R be a binary relation on it defined via $R = \{(a, a), (l, l), (n, n), (n, h), (h, n), (l, n), (g, c), (c, g), (g, l), (b, g), (g, b), (h, g), (a, b), (b, c), (h, a), (a, c)\}$. Then $\langle S, R \rangle$ is a PRAS.

If P is the reflexive closure of R (that is $P = R \cup \Delta_S$), then $\langle S, P \rangle$ is a PRAX. The successor neighborhoods associated with different elements of S are as follows (**E** is a variable taking values in S):

E	a	b	c	g	e	f	h	l	n
$[E]$	$\{a, h\}$	$\{b, c, g\}$	$\{b, c, g\}$	$\{b, c, g, h\}$	$\{e\}$	$\{f\}$	$\{h, n\}$	$\{l, g\}$	$\{n, l, g, h\}$
$[E]_o$	$\{a\}$	$\{b, c, g\}$	$\{b, c, g\}$	$\{b, c, g\}$	$\{e\}$	$\{f\}$	$\{h, n\}$	$\{l\}$	$\{n, h\}$

If $A = \{a, h, f\}$, then $A^l = \{a, h, f\}$, while $A^{lo} = \{a, f\}$ and $A^{lo} \subset A^l$. If $F = \{l\}$, then $F^l = \emptyset$, while $F^{lo} = F$ and $F^l \subset F^{lo}$.

Now let $Z = N \cup S \cup X$, where N is the set of naturals, X is the set of elements of the infinite sequences $\{x_i\}$, $\{y_j\}$. Let Q be a relation on Z such that $Q \cap S^2 = P$, $Q \cap N^2$ be some equivalence. Further let $(\forall i \in N)(i, x_{3i+1}), (x_{2i}, i), (x_i, x_{i+1}), (y_i, y_{i+1}) \in Q$. For any $i \in N$, let $P_i = \{y_k : k \neq 2j \& k < i\} \cup \{x_{2j} : 2j < i\}$ - this will be used in later sections. The extension of the example to involve nets and densely ordered subsets is standard.

Caste Hierarchy Databases:

In detailed NSS datasets on the evil interaction between castes in India, let $\mathfrak{U}_\alpha xy$ denote the state that 'members of caste x ' regard 'members of caste y ' as *untouchable* for interaction α . For suitable subsets of α , the relation is proto-transitive and reflexive. Semantics of the associated PRAX can help in deciding on social reform strategies.

3 Approximations and Definite Elements in a PRAX

Definition 2. By a Proto Approximation Space S (PRAS for short), we will mean a pair of the form $\langle \underline{S}, R \rangle$ with $\underline{S}$ being a set and R being a proto-transitive

relation on it. If R is also reflexive, then it will be called a Reflexive Proto Approximation Space (PRAX) for short). $\underline{S}$ may be infinite.

If S is a PRAX, then we will respectively denote successor and symmetrized successor neighborhoods generated by an element $x \in S$ $[x] = \{y; Ryx\}$ and $[x]_o = \{y; Ryx \& Rxy\}$ respectively. Taking these as granules, the associated granulations will be denoted by $\mathcal{G} = \{[x] : x \in S\}$ and $\mathcal{G}_o$ respectively. In all that follows S will be a PRAX unless indicated otherwise.

Definition 3. *Definable approximations on S include ($A \subseteq S$):*

Upper Proto: $A^u = \bigcup_{[x] \cap A \neq \emptyset} [x]$.

Lower Proto: $A^l = \bigcup_{[x] \subseteq A} [x]$.

Symmetrized Upper Proto $A^{uo} = \bigcup_{[x]_o \cap A \neq \emptyset} [x]_o$.

Symmetrized Lower Proto $A^{lo} = \bigcup_{[x]_o \subseteq A} [x]_o$.

Point-wise Upper $A^{u+} = \{x : [x] \cap A \neq \emptyset\}$.

Point-wise Lower $A^{l+} = \{x : [x] \subseteq A\}$.

Proposition 2. *For any subset $A \subseteq S$, all of the following hold:*

- ★ It is possible that $A^l \neq A^{l+}$ and in general, $A^l \parallel A^{lo}$.
- ★ $A \subseteq S$, $A^{uo} \subseteq A^u$.

Proof. Clearly $(\forall x) [x]_o \subseteq [x]$, but the first example includes a counter example for $A^l \parallel A^{lo}$.

Since $[x]_o \cap A \neq \emptyset$, therefore $A^{uo} = \bigcup_{[x]_o \cap A \neq \emptyset} [x]_o \subseteq \bigcup_{[x] \cap A \neq \emptyset} [x]_o \subseteq A^u = \bigcup_{[x] \cap A \neq \emptyset} [x] = A^u$. $\square$

Definition 4. *If X is an approximation operator, then by a X -definite element, we will mean a subset A satisfying $A^X = A$. The set of all X -definite elements will be denoted by $\delta_X(S)$, while the set of X and Y -definite elements (Y being another approximation operator) will be denoted by $\delta_{XY}(S)$. In particular, we will speak of lower proto-definite, upper proto definite and proto-definite elements (those that are both lower and upper proto-definite).*

Proposition 3. $\delta_u(S)$ is a complete sublattice of $\wp(S)$ with respect to inclusion.

Proof. As R is reflexive, if A, B are upper proto definite, then $A \cup B$ and $A \cap B$ are both upper proto definite. So the result holds. $\square$

Theorem 1. *In a PRAX S , the following hold:*

- ★ $\delta_u(S) \subseteq \delta_{uo}(S)$, but $\delta_{lo}(S) = \delta_{uo}(S)$.
- ★ $\delta_l(S) \parallel \delta_{lo}(S)$ in general.
- ★ It is possible that $\delta_u \not\subseteq \delta_{uo}$.

Proof. If $A \in \delta_u$, then $(\forall x \in A) [x] \subseteq A$ and $(\forall x \in A^c) [x] \cap A = \emptyset$.

So $(\forall x \in A^c) [x]_o \cap A = \emptyset$. But as $A \subseteq A^{uo}$ is necessary, we must have $A \in \delta_{uo}$. $\square$

A^{u+} , A^{l+} like operators have been more commonly used in the literature and the only kind of approximation studied in [10] for example.

Theorem 2. *All of the following hold in PRAX A:*

- ★ $(\forall x) x^{cl+} = x^{u+c}$, $x^{cu+} = x^{l+c}$ - that is $l+$ and $u+$ are mutually dual
- ★ $u+$ ($l+$ resp.) is a monotone $\vee$ - ($\text{complete } \wedge$ - resp.) morphism.
- ★ $\partial(x) = \partial(x^c)$, where ∂ stands for the boundary operator.
- ★ $\Im(u+)$ is an interior system while $\Im(l+)$ is a closure system.
- ★ $\Im(u+)$ and $\Im(l+)$ are dually isomorphic lattices.

Theorem 3. $(\forall A \in \wp(S)) A^{l+} \subseteq A^l$, $A^{u+} \subseteq A^u$.

Proof. ★ If $x \in A^{l+}$, then $[x] \subseteq A$ and so $[x] \subseteq A^l$, $x \in A^l$.

★ If $x \in A^l$, then $(\exists y \in A)[y] \subseteq A$, Rxy . But it is possible that $[x] \not\subseteq A$, therefore it is possible that $x \notin A^{l+}$ and $A^l \not\subseteq A^{l+}$.

★ If $x \in A^{u+}$, then $[x] \cap A \neq \emptyset$, so $x \in A^u$.

★ So $A^{u+} \subseteq A^u$.

★ Note that $x \in A^u$, if and only if $(\exists z \in S) x \in [z]$, $[z] \cap A \neq \emptyset$, but this does not imply $x \in A^{u+}$.

Theorem 4. *In a PRAX, all of the following hold:*

Bi $(\forall A \in \wp(S)) A^{ll} = A^l \& A^u \subseteq A^{uu}$.

l-Cup $(\forall A, B \in \wp(S)) A^l \cup B^l \subseteq (A \cup B)^l$.

l-Cap $(\forall A, B \in \wp(S)) (A \cap B)^l \subseteq A^l \cap B^l$.

u-Cup $(\forall A, B \in \wp(S)) (A \cup B)^u = A^u \cup B^u$

u-Cap $(\forall A, B \in \wp(S)) (A \cap B)^u \subseteq A^u \cap B^u$

Dual $(\forall A \in \wp(S)) A^{lc} \subseteq A^{cu}$.

Proof. **l-Cup** For any $A, B \in \wp S$, $x \in (A \cup B)^l$

$\Leftrightarrow (\exists y \in (A \cup B)) x \in [y] \subseteq A \cup B$.

$\Leftrightarrow (\exists y \in A) x \in [y] \subseteq A \cup B$ or $(\exists y \in B) x \in [y] \subseteq A \cup B$.

$\Leftrightarrow (\exists y \in A) x \in [y] \subseteq A$ or $(\exists y \in A) x \in [y] \subseteq B$ or $(\exists y \in B) x \in [y] \subseteq A$ or $(\exists y \in B) x \in [y] \subseteq B$ - this is implied by $x \in A^l \cup B^l$.

l-Cap For any $A, B \in \wp S$, $x \in (A \cap B)^l$

$\Leftrightarrow x \in A \cap B$

$\Leftrightarrow (\exists y \in A \cap B) x \in [y] \subseteq A \cap B$ and $x \in A$, $x \in B$

$\Leftrightarrow (\exists y \in A) x \in [y] \subseteq A$ and $(\exists y \in B) x \in [y] \subseteq B$ - Clearly this statement implies $x \in A^l \& x \in B^l$, but the converse is not true in general.

u-Cup $x \in (A \cup B)^u$

$\Leftrightarrow x \in \bigcup_{[y] \cap (A \cup B) \neq \emptyset} [y]$

$\Leftrightarrow x \in \bigcup_{([y] \cap A) \cup ([y] \cap B) \neq \emptyset} [y]$

$\Leftrightarrow x \in \bigcup_{[y] \cap A \neq \emptyset} [y]$ or $x \in \bigcup_{[y] \cap B \neq \emptyset} [y]$

$\Leftrightarrow x \in A^u \cup B^u$.

u-Cap By monotonicity, $(A \cap B) \subseteq A^u$ and $(A \cap B) \subseteq B^u$, so $(A \cap B)^u \subseteq A^u \cap B^u$.


Dual If $z \in A^{lc}$, then $z \in [x]^c$ for all $[x] \subseteq A$ and either, $z \in A \setminus A^l$ or $z \in A^c$.
 If $z \in A^c$ then $z \in A^{cu}$. If $z \in A \setminus A^l$ and $z \neq A^{cu} \setminus A^c$ then $[z] \cap A^c = \emptyset$. But this contradicts $z \notin A^{cu} \setminus A^c$. So $(\forall A \in \wp(S)) A^{lc} \subseteq A^{cu}$.

Theorem 5. In a PRAX S , all of the following hold:

1. $(\forall A, B \in \wp(S)) (A \cap B)^{l+} = A^{l+} \cap B^{l+}$
2. $(\forall A, B \in \wp(S)) A^{l+} \cup B^{l+} \subseteq (A \cup B)^{l+}$
3. $(\forall A \in \wp(S)) (A^{l+})^c = (A^c)^{u+}$, & $A^{l+} \subseteq A^{lo}$ & $A^{uo} \subseteq A^{u+}$ & $A^{l+} \subseteq A^{lo}$.

Proof. 1. $x \in (A \cap B)^{l+} \Leftrightarrow [x] \subseteq A \cap B \Leftrightarrow [x] \subseteq A$ and $[x] \subseteq B \Leftrightarrow x \in A^{l+}$ and $x \in B^{l+}$.
 2. $x \in A^{l+} \cup B^{l+} \Leftrightarrow [x] \subseteq A^{l+}$ or $[x] \subseteq B^{l+} \Leftrightarrow [x] \subseteq A$ or $[x] \subseteq B \Rightarrow [x] \subseteq A \cup B \Leftrightarrow x \in (A \cup B)^{l+}$.
 3. $z \in A^{l+} \Leftrightarrow z \notin A^c \Leftrightarrow [z] \not\subseteq A^c \Leftrightarrow z \cap A^c \neq \emptyset$.
 If $x \in A^{l+}$, then $[x] \subseteq A$. But as $[x]_o \subseteq [x]$, $A^{l+} \subseteq A^{lo}$. □

From the above, we have the following relation between approximations in general ($u+ \longrightarrow u$ should be read as *the $u+$ -approximation of a set is included in the u -approximation of the same set*):



If a relation R is purely reflexive and not proto-transitive on a set S , then the relation $\tau(R) = R \cap R^{-1}$ will not be an equivalence and for a $A \subset S$, it is possible that $A^{uol} \subseteq A$ or $A^{uol} \parallel A$ or $A \subseteq A^{uol}$.

3.1 Algebras of Rough Definite Elements

Theorem 6. On the set of proto definite elements $\delta_{lu}(S)$ of a PRAX S , we can define the following:

1. $x \wedge y \stackrel{\Delta}{=} x \cap y$; $x \vee y \stackrel{\Delta}{=} x \cup y$
2. $0 \stackrel{\Delta}{=} \emptyset$; $1 \stackrel{\Delta}{=} S$; $x^c \stackrel{\Delta}{=} S \setminus x$

The resulting algebra $\delta_{proto}(S) = \langle \delta_{lu}(S), \vee, \wedge, c, 0, 1 \rangle$ is a Boolean lattice.

Proof. We need to show that the operations are well defined. Suppose x, y are proto-definite elements, then

1. $(x \cap y)^u \subseteq x^u \cap y^u = x \cap y$. But, $(x \cap y)^l = (x^u \cap y^u)^l = (x \cap y)^{ul} = (x \cap y)^u = x \cap y$. Since $a^{ul} = a^u$ for any a . Further, $(x \cup y)^u = x \cup y = x^l \cup y^l \subseteq (x \cup y)^l$.
2. Suppose $A \in \delta_{lu}(S)$, then $(\forall z \in A^c) [z] \cap A = \emptyset$ is essential, else $[z]$ would be in A^u . This means $[z] \subseteq A^c$ and so $A^c = A^{cl}$. If there exists a $a \in A$ such that $[a] \cap A^c \neq \emptyset$, then $[a] \subseteq A^u = A$. So $A^c \in \delta_{lu}(S)$.

The lattice order can be defined via, $x \leq y$ if and only if $x \cup y = y$ and $x \cap y = x$. □

4 The Representation of Roughly Equal Elements

The representation of requal elements in terms of definite elements are well known in case classical rough set theory. In case of more general spaces including tolerance spaces [1], most authors have been concerned with describing the interaction of rough approximations of different types and not of the interaction of requal objects. Higher order approaches, developed by the present author as in [4] for bitten approximation spaces, permit constructs over sets of requal objects. In the light of the contamination problem [4,1], it would be an improvement to describe without higher order constructs. In this section a new method of representing requal elements based on expanding concepts of definite elements is developed.

Definition 5. A subset A of $\wp(S)$ will be said to a set of requal elements if and only if $(\forall x, y \in A) x^l = y^l \& x^u = y^u$. It will be said to be full if no other subset properly including A has the property.

Relative the situation for a general RYS, we have

Theorem 7 (Meta-Theorem). In a PRAX S , full set of requal elements is necessarily a union of intervals in $\wp(S)$.

Definition 6. A non-empty set of non singleton subsets $\alpha = \{x : x \subseteq \wp(S)\}$ will be said to be a upper broom if and only if all of the following hold:

- ★ $(\forall x, y \in \alpha) x^u = y^u \& x \parallel y$
- ★ If $\alpha \subset \beta$, then β fails to satisfy at least one of the two conditions.

Definition 7. A non-empty set of non singleton subsets $\alpha = \{x : x \subseteq \wp(S)\}$ will be said to be a lower broom if and only if all of the following hold:

- ★ $(\forall x, y \in \alpha) x^l = y^l \neq x$
- ★ $(\forall x, y \in \alpha) x \parallel y$
- ★ If $\beta \subset \alpha$ and β has at least two elements, then β fails to satisfy at least one of the two conditions.

$\psi(S), \cap(S)$ will respectively denote the set of lower and upper brooms of S .

Proposition 4. If $x \in \delta_{lu}(S)$ then $\{x\} \notin \cap(S)$ and $\{x\} \notin \psi(S)$.

In the next definition, the concept of union of intervals in a partially ordered set is modified in a way for use with specific types of objects.

Definition 8. By a bruinval, we will mean a subset of $\wp(S)$ of one of the following forms $\alpha, \beta \in \wp(\wp(S))$:

- ★ Bruinval-0: Intervals of the form $(x, y), [x, y), [x, x], (x, y]$ for $x, y \in \wp(S)$.
- ★ Open Bruinvals: $[x, \alpha) = \{z : x \leq z < b \& b \in \alpha\}, (x, \alpha] = \{z : x < z \leq b \& b \in \alpha\}$ and $(x, \alpha) = \{z : x < z < b, b \in \alpha\}$.
- ★ Closed Bruinvals: $[x, \alpha] = \{z : x \leq z \leq b \& b \in \alpha\}$.

- ★ *Closed Set Bruinvals:* $[\alpha, \beta] = \{z : x \leq z \leq y \text{ \& } x \in \alpha \text{ \& } y \in \beta\}$.
- ★ *Open Set Bruinvals:* $(\alpha, \beta) = \{z : x < z < y, x \in \alpha \text{ \& } y \in \beta\}$.
- ★ *Semi-Closed Set Bruinvals:* $[[\alpha, \beta]]$ defined as follows: $\alpha = \alpha_1 \cup \alpha_2, \beta = \beta_1 \cup \beta_2$
and $[[\alpha, \beta]] = (\alpha_1, \beta_1) \cup [\alpha_2, \beta_2] \cup (\alpha_1, \beta_2] \cup [\alpha_2, \beta_1)$.

In the example of the second section, the representation of the rough object (P_i^l, P_i^u) requires set bruinvals.

Theorem 8. *A bruinval-0 of the form (x, y) is a full set of requal elements if $x, y \in \delta_{lu}(S)$ and x is covered by y in the order on $\delta_{lu}(S)$.*

If $x, y \in \delta_{lu}(S)$ then sets of the form $[x, y), (x, y]$ cannot be a non-empty set of requal elements, while those of the form $[x, y]$ can be if and only if $x = y$.

Proposition 5. *An interval $[x, y)$ is a full set of requal elements if*

- ★ $x^l, y^u \in \delta_{lu}(S), x^l = y^l$ and $x^u = y^u$,
- ★ x^l is covered by y^u in $\delta_{lu}(S)$ and
- ★ $x \setminus (x^l)$ and $y^u \setminus y$ are singletons

In the above proposition the condition $x^l, y^u \in \delta_{lu}(S)$, is not necessary.

Theorem 9. *If a bruinval-0 of the form $[x, y]$ satisfies*

- ★ $x^l = y^l = x$ and $x^u = y^u$,
- ★ $y^u \setminus y$ is a singleton.

then $[x, y]$ is a full set of requal objects.

Proof. Under the conditions, if $[x, y]$ is not a full set of requal objects, then there must exist at least one set h such that $h^l = x$ and $h^u = y^u$ and $h \notin [x, y]$. But this contradicts the order constraint $x^l \leq h y^u$. Note that $y^u \notin [x, y]$ under the conditions. $\square$

Theorem 10. *If a bruinval-0 of the form $(x, y]$ satisfies*

- ★ $x^l = y^l = x$ and $(\forall z \in (x, y]) z^u = y^u$,
- ★ $y^u \setminus y$ is a singleton.

then $(x, y]$ is a full set of requal objects, that does not intersect the full set $[x, x^u]$.

Proof. By monotonicity it follows that $(x, y]$ is a full set of requal objects. then there must exist at least one set h such that $h^l = x$ and $h^u = y^u$ and $h \notin [x, y]$. But this contradicts the order constraint $x^l \leq h y^u$. Note that $y^u \notin [x, y]$ under the conditions. $\square$

Theorem 11. *A bruinval-0 of the form (x^l, x^u) is not always a set of requal elements, but will be so when $x^{uu} = x^u$. In the latter situation it will be full if $[x^l, x^u)$ is not full.*

Most of the above considerations extend to other types of bruinvals as is shown below and remain amenable.

Theorem 12. *Bruinval (x, α) is a full set of requal elements if and only if α is an upper broom, $(\forall y \in \alpha) x^l = y^l, x^u = y^u$ and $(\forall z)(x^l \subseteq z \subseteq x \longrightarrow z^u \subseteq x^u)$.*

Proof. It is clear that for any $y \in \alpha$, (x, y) is a convex interval and all elements in it have same upper and lower approximations. The third condition ensures that $[z, \alpha)$ is not a full set for any $z \in [x^l, x)$. $\square$

Definition 9. *An element $x \in \wp(S)$ will be said to be a weak upper critical element relative $z \subset x$ if and only if $(\forall y \in \wp(S))(z = y^l \& x \subset y \longrightarrow x^u \subset y^u)$.*

An element $x \in \wp(S)$ will be said to be an upper critical element relative $z \subset x$ if and only if $(\forall v, y \in \wp(S))(z = y^l = v^l \& v \subset x \subset y \longrightarrow v^u = x^u \subset y^u)$. Note that the inclusion is strict.

An element a will be said to be bicritical relative b if and only if $(\forall x, y \in \wp(S))(a \subset x \subseteq y \subset b \longrightarrow x^u = y^u \& x^l = y^l \& x^u \subset b^u \& a^l \subset x^l)$.

If x is an upper critical point relative z , then $[z, x)$ or (z, x) is a set of roughly equivalent elements.

Definition 10. *An element $x \in \wp(S)$ will be said to be an weak lower critical element relative $z \supset x$ if and only if $(\forall y \in \wp(S))(z = y^u \& y \subset x \longrightarrow y^l \subset x^l)$.*

An element $x \in \wp(S)$ will be said to be an lower critical element relative $z \supset x$ if and only if $(\forall y, v \in \wp(S))(z = y^u = v^u \& y \subset x \subset v \longrightarrow y^l \subset x^l = v^l)$.

An element $x \in \wp(S)$ will be said to be an lower critical element if and only if $(\forall y \in \wp(S))(y \subset x \longrightarrow y^l \subset x^l)$. An element that is both lower and upper critical will be said to be critical.

Proposition 6. *In a PRAX, every upper definite subset is also upper critical, but the converse need not hold.*

The most important thing about the different lower and upper critical points is that they help in determining full sets of requal elements by determining the boundaries of intervals in bruinvals of different types.

4.1 Atoms in the POSET of Roughly Equivalent Objects

Proposition 7. *For $x, y \in \wp(S) | \approx = H$ (say), if we define $x \leq y$ if and only if $(\forall a \in x)(\forall b \in y)a^l \subseteq b^l \& a^u \subseteq b^u$, then $\leq$ is a bounded and directed partial order. The least element will be denoted by 0 ($0 = \{\emptyset\}$) and the greatest by 1 ($1 = \{S\}$). An atom of H is any element that covers 0. The set of atoms of H will be denoted by $At(H)$.*

Theorem 13. *H is atomic and atoms of H will be bruinvals of specific types:*

Type-0: $(\emptyset, [x])$ that intersect no other set of requal sets.

Type-1: $(\emptyset, \alpha)$ that do not contain full sets of requal sets.

Type-2: (α, β) not containing full sets of requal sets and s.t. $(\forall x)x^l = \emptyset$.

Proof. That H is atomic will be proved last. It is obvious that a bruinval of the form (α, β) can be an atom only if α is the $\emptyset$. If not, then each element x of the bruinval $(\emptyset, \alpha)$ will satisfy $x^l = \emptyset \subset x^u$, thereby contradicting the assumption that (α, β) is an atom.

If $[x]$ intersects no other successor neighborhood, then $(\forall y \in (\emptyset, [x]))y^l = \emptyset \& x^u = [x]$ and it will be a minimal set of requal elements containing 0. The other part can be verified based on the representation of possible sets of roughly equivalent elements.

To prove H is atomic, we need to show that any element x greater than 0 is either an atom or there exists an atom a such that $a \leq x$, that is $(\forall x)(\exists a \in At(H))(0 < x \longrightarrow a \leq x)$.

Suppose the bruinval (α, β) represents a non-atom, then it is necessary that $(\forall x \in \alpha)x^l \neq \emptyset \& x^u \subseteq S$.

Suppose the neighborhoods included in x^u are $\{[y] : y \in B \subseteq S\}$. If all combinations of bruinvals of the form $(\emptyset, \gamma)$ formed from these neighborhoods are not atoms, then it is necessary that the upper approximation of every singleton subset of a set in γ properly contains another non-trivial upper approximation. This is impossible and so H must be atomic. $\square$

Theorem 14. *The granulation $\mathcal{G}$ satisfies RA, LCG, MER, ST [1] with respect to the approximations l, u , but $\mathcal{G} \cup \mathcal{G}_o$ fails LCG for the approximations l, u, l_o, u_o .*

5 Geometry of Granular Knowledge Interpretation

In my opinion, *Any knowledge, however involved, may be seen as a collection of concepts with admissible operations of reasoning defined on them.* Knowledges associated PRAX have various peculiarities corresponding to the semantic evolution of rough objects in it. The semantic domains of representation properly contain the semantic domains of interpretation. Not surprisingly, it is because the rough objects corresponding to l, u cannot be represented perfectly in terms of objects from $\delta_{lu}(S)$ alone. *In the nongranular perspective too, this representation aspect should matter - "should",* because it is matter of choice during generalization from the classical case in the non granular approach.

The natural rough semantic domains of l, u is Meta-R, while that of l_o, u_o is $\mathfrak{D}$ (say, corresponding rough objects of $\tau(R)$). These can be seen as separate domains or as parts of a minimal containing domain that permits enough expression. The main problem of granular KI is that knowledge is correctly representable in terms of *atomic concepts of knowledge* at semantic domains placed between Meta-C and Meta-R and not at the latter. So the characterization of possible semantic domains and their mutual ordering - leading to their geometry is of interest.

The following will be assumed to be *part* of the interpretation:

- ★ Two types of rough objects corresponding to Meta-R and $\mathfrak{D}$ and their natural correspondence correspond to concepts or weakenings thereof. A concept relative one semantic domain need not be one of the other.

- ★ A granule of the rough semantic domain $\mathcal{O}$ is necessarily a concept of $\mathcal{O}$, but a granule of Meta-R may not be a concept of $\mathcal{O}$ or Meta-R.
- ★ Critical points are not necessarily concepts of either semantic domain.
- ★ Critical points and the representation of rough objects require the rough semantic domains to be extended.

The above obviously assumes that a PRAX S has at least two kinds of knowledge associated (in relation to the Pawlak-sense interpretation). To make the interpretations precise, we will indicate them by $\mathcal{I}_1(S)$ and $\mathcal{I}_o(S)$ respectively (corresponding to the approximations to l , u and l_o , u_o respectively). The pair $(\mathcal{I}_1(S), \mathcal{I}_o(S))$ will also be referred to as the *generalized KI*.

Definition 11. *Given two PRAX $S = \langle \underline{S}, R \rangle$, $V = \langle \underline{S}, Q \rangle$, S will be said to be o-coarser than V if and only if $\mathcal{I}_o(S)$ is coarser than $\mathcal{I}_o(V)$ in Pawlak-sense (that is $\tau(R) \subseteq \tau(Q)$). Conversely, V will be said to be a o-refinement of S .*

S will be said to be p-coarser than V if and only if $\mathcal{I}_1(S)$ is coarser than $\mathcal{I}_1(V)$ in the sense $R \subseteq Q$. Conversely, V will be said to be a p-refinement of S .

An extended concept of positive regions is defined next.

Definition 12. *If $S_1 = \langle \underline{S}, Q \rangle$ and $S_2 = \langle \underline{S}, P \rangle$ are two PRAX such that $Q \subset R$, then by the granular positive region of Q with respect to R is given by $gPOS_R(Q) = \{[x]_Q^{l_R} : x \in \underline{S}\}$, where $[x]_Q^{l_R}$ is the lower approximation (relative R) of the Q -related elements of x . Using this we can define the granular extent of dependence of knowledge encoded by R on the knowledge encoded by Q by natural injections : $gPOS_R(Q) \mapsto \mathcal{G}_R$.*

Lower critical points can be naturally interpreted as precepts that are definitely included in the discourse, while upper critical points are precepts that include most of the discourse. The problem with this interpretation is that it's representation requires a semantic domain at which critical points of different kinds can be found. A key requirement for such a domain would be the feasibility of rough counting procedures like IPC [1]. We will refer to a semantic domain that has critical points of different types as basic objects as a **Meta-RC**.

The following possible axioms of granular knowledge that also figure in my earlier paper [2], get into difficulties with the present approach and even when we restrict attention to $\mathcal{I}_1(S)$:

1. Individual granules are atomic units of knowledge.
2. Maximal collections of granules subject to a concept of mutual independence are admissible concepts of knowledge.
3. Parts common to subcollections of maximal collections of granules are also knowledge.

The first axiom holds in weakened form as the granulation $\mathcal{G}$ for $\mathcal{I}_1(S)$ is only lower definite and affects the other. The possibility of other nice granulations being possible for the PRAX case appears to be possible at the cost of other nice properties (14). So we can conclude that in proper KR happens at semantic domains like **Meta-RC** where critical points of different types are perceived.

Further at Meta-R, rough objects may correspond to knowledge or conjectures - if we require the concept of proof to be an ontological concept or beliefs. The scenario can be made more complex with associations of $\mathfrak{D}$ knowledges.

From a non-granular perspective, in Meta-R rough objects must correspond to knowledge with some of them lacking a proper evolution - there is no problem here. Even if we permit $\mathfrak{D}$ objects, then in the perspective we would be able to speak of two kinds of closely associated knowledges.

Concluding Remarks:

In this research we have developed a new general rough set theory over proto transitive relations, the representation of rough objects and definite objects. The knowledge interpretation of rough sets is also generalized to provide sensible knowledge interpretations across different semantic domains. This paves the way for possible semantics, measures and/or logics of knowledge consistency. We have also shown in concrete terms that granular KI and classical KI are different things.

References

1. Mani, A.: Dialectics of Counting and the Mathematics of Vagueness. Transactions on Rough Sets **XV**(LNCS 7255) (2012) 122–180
2. Mani, A.: Towards Logics of Some Rough Perspectives of Knowledge. In Suraj, Z., Skowron, A., eds.: Intelligent Systems Reference Library dedicated to the memory of Prof. Pawlak,. Springer Verlag (2011-12) 342–367
3. Mani, A.: Granular Approach to Generalised Transitive Rough Sets. In Process (2012) 1–87
4. Mani, A.: Choice Inclusive General Rough Semantics. Information Sciences **181**(6) (2011) 1097–1115
5. Chajda, I., Haviar, M.: Induced Pseudo Orders. Acta Univ. Palack. Olomou **30**(1) (1991) 9–16
6. Mani, A.: Esoteric Rough Set Theory-Algebraic Semantics of a Generalized VPRS and VPRFS. In Skowron, A., Peters, J.F., eds.: Transactions on Rough Sets VIII. Volume LNCS 5084. Springer Verlag (2008) 182–231
7. Mani, A.: Axiomatic Approach to Granular Correspondences. In Li, T., et al., eds.: Proceedings of RSKT'2012, LNAI 7414. Volume LNAI 7414., Springer-Verlag (2012) 482–487
8. Cattaneo, G., Ciucci, D.: Lattices with Interior and Closure Operators and Abstract Approximation Spaces. In Peters, J.F., et al., eds.: Transactions on Rough Sets X, LNCS 5656. Springer (2009) 67–116
9. Ciucci, D.: Approximation Algebra and Framework. Fundamenta Informaticae **94** (2009) 147–161
10. Jarvinen, J.: Lattice Theory for Rough Sets. In Peters, J.F., et al., eds.: Transactions on Rough Sets VI. Volume LNCS 4374. Springer Verlag (2007) 400–498